arXiv:1005.0455v1 [math.GM] 4 May 2010

ON THE WEIGHTED OSTROWSKI TYPE INTEGRAL INEQUALITY FOR DOUBLE INTEGRALS

MEHMET ZEKI SARIKAYA AND HASAN OGUNMEZ

Abstract. In this paper, we establish new an inequality of weighted Ostrowski-type for double integrals involving functions of two independent variables by using fairly elementary analysis.

1. Introduction

In 1938, the classical integral inequality established by Ostrowski [3] as follows:

Theorem 1. *Let* $f:[a,b]\rightarrow\mathbb{R}$ *be a differentiable mapping on* (a,b) *whose derivative* $f^{'}:(a,b)\rightarrow\mathbb{R}$ *is bounded on* (a,b)*, i.e.,* $\|f'\|_{\infty}=\sup_{t\in(a,b)}|f'(t)|<\infty$*. Then, the inequality holds:*

$$\left|f(x)-\frac{1}{b-a}\int_{a}^{b}f(t)dt\right|\leq\left[\frac{1}{4}+\frac{(x-\frac{a+b}{2})^2}{(b-a)^2}\right](b-a)\|f'\|_{\infty}\tag{1.1}$$

for all $x\in[a,b]$*. The constant* $\frac{1}{4}$ *is the best possible.*

In a recent paper [1], Barnett and Dragomir proved the following Ostrowski type inequality for double integrals

Theorem 2. *Let* $f:[a,b]\times[c,d]\rightarrow\mathbb{R}$ *be continuous on* $[a,b]\times[c,d]$*,* $f''_{x,y}=\frac{\partial^2 f}{\partial x\partial y}$ *exists on* $(a,b)\times(c,d)$ *and is bounded, i.e.,*

$$\left\|f''_{x,y}\right\|_{\infty}=\sup_{(x,y)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f(x,y)}{\partial x\partial y}\right|<\infty.$$

Then, we have the inequality:

$$\begin{aligned}&\left|\int_{a}^{b}\int_{c}^{d}f(s,t)dtds-(d-c)(b-a)f(x,y)\right.\\&\qquad\left.-\left[(b-a)\int_{c}^{d}f(x,t)dt+(d-c)\int_{a}^{b}f(s,y)ds\right]\right|\\&\leq\left[\frac{1}{4}(b-a)^2+(x-\frac{a+b}{2})^2\right]\left[\frac{1}{4}(d-c)^2+(y-\frac{d+c}{2})^2\right]\left\|f''_{x,y}\right\|_{\infty}\end{aligned}\tag{1.2}$$

for all $(x,y)\in[a,b]\times[c,d]$*.*

2000 *Mathematics Subject Classification.* 26D07, 26D15.

Key words and phrases. Ostrowski's inequality.

In [1], the inequality (1.2) is established by the use of integral identity involving Peano kernels. In [5], Pachpatte obtained an inequality in the view (1.2) by using elementary analysis. The interested reader is also refered to ([1], [2], [4]-[8]) for Ostrowski type inequalities in several independent variables.

The main aim of this note is to establish a new weighted Ostrowski type inequality for double integrals involving functions of two independent variables and their partial derivatives.

2. Main Result

Throughout this work, we assume that the weight function $w:[a,b]\to[0,\infty)$, is integrable, nonnegative and

$$m(a,b)=\int_a^b w(t)dt<\infty. \tag{2.1}$$

Lemma 1. *Let $f:[a,b]\times[c,d]\to\mathbb{R}$ be an absolutely continuous fuction such that the partial derivative of order 2 exist for all $(t,s)\in[a,b]\times[c,d]$. Then, we have*

$$\begin{aligned} f(x,y) &= \frac{1}{m(a,b)}\int_a^b w(t)f(t,y)dt+\frac{1}{m(c,d)}\int_c^d w(s)f(x,s)ds \\ &\quad (2.2)-\frac{1}{m(a,b)m(c,d)}\left[\int_a^b\int_c^d w(t)w(s)f(t,s)dsdt-\int_a^b\int_c^d p(x,t)q(y,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt\right]\end{aligned}$$

where

$$p(x,t)=\begin{cases} p_1(a,t)=\int_a^t w(u)du, & a\le t<x \\ p_2(b,t)=\int_b^t w(u)du, & x\le t\le b\end{cases}$$

and

$$q(y,s)=\begin{cases} q_1(c,s)=\int_c^s w(u)du, & c\le s<y \\ q_2(d,s)=\int_d^s w(u)du, & y\le s\le d.\end{cases}$$

Proof. By definitions of $p(x,t)$ and $q(y,s)$, we have

$$\begin{aligned}
\int_a^b\int_c^d p(x,t)q(y,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt &= \int_a^x\int_c^y p_1(a,t)q_1(c,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt \\
&+\int_a^x\int_y^d p_1(a,t)q_2(d,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt+\int_x^b\int_c^y p_2(b,t)q_1(c,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt \\
&+\int_x^b\int_y^d p_2(b,t)q_2(d,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt.
\end{aligned}\tag{2.3}$$

Integrating by parts, we can state:

$$\begin{aligned}
&\int_a^x\int_c^y p_1(a,t)q_1(c,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt \\
&=\int_a^x p_1(a,t)\left[q_1(c,y)\frac{\partial f(t,y)}{\partial t}-\int_c^y w(s)\frac{\partial f(t,s)}{\partial t}ds\right]dt \\
&=p_1(a,x)q_1(c,y)f(x,y)-q_1(c,y)\int_a^x w(t)f(t,y)dt \\
&-p_1(a,x)\int_c^y w(s)f(x,s)ds+\int_a^x\int_c^y w(s)w(t)f(t,s)dsdt.
\end{aligned}\tag{2.4}$$

$$\begin{aligned}
&\int_a^x\int_y^d p_1(a,t)q_2(d,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt \\
&=\int_a^x p_1(a,t)\left[-q_2(d,y)\frac{\partial f(t,y)}{\partial t}-\int_y^d w(s)\frac{\partial f(t,s)}{\partial t}ds\right]dt \\
&=-p_1(a,x)q_2(d,y)f(x,y)+q_2(d,y)\int_a^x w(t)f(t,y)dt \\
&-p_1(a,x)\int_y^d w(s)f(x,s)ds+\int_a^x\int_y^d w(s)w(t)f(t,s)dsdt.
\end{aligned}\tag{2.5}$$

$$\begin{aligned}
&\int_x^b\int_c^y p_2(b,t)q_1(c,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt \\
&= \int_x^b p_2(b,t)\left[q_1(c,y)\frac{\partial f(t,y)}{\partial t} - \int_c^y w(s)\frac{\partial f(t,s)}{\partial t}ds\right]dt \\
&= -p_2(b,x)q_1(c,y)f(x,y) - q_1(c,y)\int_x^b w(t)f(t,y)dt \\
&+p_2(b,x)\int_c^y w(s)f(x,s)ds + \int_x^b\int_c^y w(s)w(t)f(t,s)dsdt.
\end{aligned} \tag{2.6}$$

$$\begin{aligned}
&\int_x^b\int_y^d p_2(b,t)q_2(d,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt \\
&= \int_x^b p_2(b,t)\left[q_2(d,y)\frac{\partial f(t,y)}{\partial t} - \int_y^d w(s)\frac{\partial f(t,s)}{\partial t}ds\right]dt \\
&= p_2(b,x)q_2(d,y)f(x,y) + q_2(d,y)\int_x^b w(t)f(t,y)dt \\
&+p_2(b,x)\int_y^d w(s)f(x,s)ds + \int_x^b\int_y^d w(s)w(t)f(t,s)dsdt.
\end{aligned} \tag{2.7}$$

Adding (2.4)-(2.7) and rewriting, we easily deduce:

$$\begin{aligned}
&\int_a^b\int_c^d p(x,t)q(y,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt = m(a,b)m(c,d)f(x,y) - m(c,d)\int_a^b w(t)f(t,y)dt \\
&-m(a,b)\int_c^d w(s)f(x,s)ds + \int_a^b\int_c^d w(s)w(t)f(t,s)dsdt
\end{aligned} \tag{2.8}$$

which this completes the proof. □

Theorem 3. *Let $f:[a,b]\times[c,d]\to\mathbb{R}$ be an absolutely continuous fuction such that the partial derivative of order 2 exist and is bounded, i.e.,*

$$\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty = \sup_{(x,y)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right| < \infty$$

for all $(t,s)\in[a,b]\times[c,d]$. *Then, we have*

$$\begin{aligned}&\Bigg|f(x,y)-\left[\frac{1}{m(a,b)}\int_a^b w(t)f(t,y)dt+\frac{1}{m(c,d)}\int_c^d w(s)f(x,s)ds\right]\\ &\qquad -\frac{1}{m(a,b)m(c,d)}\int_a^b\int_c^d w(s)w(t)f(t,s)dsdt\Bigg| \qquad (2.9)\\ &\leq \frac{A(x)B(y)}{m(a,b)m(c,d)}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty\end{aligned}$$

where

$$A(x)=\int_a^x (x-u)w(u)du+\int_x^b (u-x)w(u)du$$

and

$$B(y)=\int_c^y (y-u)w(u)du+\int_y^d (u-y)w(u)du.$$

Proof. From Lemma 1 and using the properties of modulus, we observe that

$$\begin{aligned}&\Bigg|f(x,y)-\left[\frac{1}{m(a,b)}\int_a^b w(t)f(t,y)dt+\frac{1}{m(c,d)}\int_c^d w(s)f(x,s)ds\right]\\ &-\frac{1}{m(a,b)m(c,d)}\int_a^b\int_c^d w(s)w(t)f(t,s)dsdt\Bigg|\\ &\leq \frac{1}{m(a,b)m(c,d)}\int_a^b\int_c^d |p(x,t)|\,|q(y,s)|\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right|dsdt \qquad (2.10)\\ &\leq \frac{1}{m(a,b)m(c,d)}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty\int_a^b\int_c^d |p(x,t)|\,|q(y,s)|\,dsdt.\end{aligned}$$

Now, using the change of order of integration we get

$$\begin{aligned}\int_a^b |p(x,t)|\,dt &= \int_a^x |p_1(a,t)|\,dt + \int_x^b |p_2(b,t)|\,dt \\ &\le \int_a^x\int_a^t w(u)dudt + \int_x^b\int_t^b w(u)dudt \\ &= \int_a^x w(u)\int_u^x dtdu + \int_x^b w(u)\int_x^u dtdu \\ &= \int_a^x (x-u)w(u)du + \int_x^b (u-x)w(u)du \end{aligned} \tag{2.11}$$

and similarly,

$$\begin{aligned}\int_c^d |q(y,s)|\,ds &= \int_c^y |q_1(c,s)|\,ds + \int_y^d |q_2(d,s)|\,ds \\ &\le \int_c^y (y-u)w(u)du + \int_y^d (u-y)w(u)du. \end{aligned} \tag{2.12}$$

Thus, using (2.11) and (2.12) in (2.10), we obtain the inequality (2.9) and the proof is completed. □

Remark 1. *If we choose $w(u)=1$ in Theorem 3, then the inequality (2.9) reduces the inequality (1.2) which is proved by Barnett and Dragomir in* [1].

Corollary 1. *Under the assumptions of Theorem 3, we have*

$$\begin{aligned}&\left| f(\frac{a+b}{2},\frac{c+d}{2}) - \left[\frac{1}{m(a,b)}\int_a^b w(t)f(t,\frac{c+d}{2})dt + \frac{1}{m(c,d)}\int_c^d w(s)f(\frac{a+b}{2},s)ds\right]\right. \\ &\qquad\left. -\frac{1}{m(a,b)m(c,d)}\int_a^b\int_c^d w(s)w(t)f(t,s)dsdt\right| \\ &\qquad\qquad \le \frac{A(\frac{a+b}{2})B(\frac{c+d}{2})}{m(a,b)m(c,d)}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty\end{aligned}$$

where

$$A(\frac{a+b}{2}) = \int_{a}^{\frac{a+b}{2}} (\frac{a+b}{2} - u)w(u)du + \int_{\frac{a+b}{2}}^{b} (u - \frac{a+b}{2})w(u)du$$

and

$$B(\frac{c+d}{2}) = \int_{c}^{\frac{c+d}{2}} (\frac{c+d}{2} - u)w(u)du + \int_{\frac{c+d}{2}}^{d} (u - \frac{c+d}{2})w(u)du.$$

Remark 2. *We choose $w(u) = 1$ in Corollary 1, we get*

$$\left| f(\frac{a+b}{2}, \frac{c+d}{2}) - \left[\frac{1}{b-a} \int_{a}^{b} f(t, \frac{c+d}{2})dt + \frac{1}{d-c} \int_{c}^{d} f(\frac{a+b}{2}, s)ds \right] - \frac{1}{(b-a)(d-c)} \int_{a}^{b}\int_{c}^{d} f(t,s)dsdt \right| \leq \frac{(b-a)(d-c)}{16} \left\| \frac{\partial^2 f(t,s)}{\partial t \partial s} \right\|_{\infty}.$$

Remark 3. *We choose $w(u) = u$ in Corollary 1, we get*

$$\left| f(\frac{a+b}{2}, \frac{c+d}{2}) - \left[\frac{2}{(b-a)^2} \int_{a}^{b} tf(t, \frac{c+d}{2})dt + \frac{2}{(d-c)^2} \int_{c}^{d} sf(\frac{a+b}{2}, s)ds \right] - \frac{4}{(b-a)^2(d-c)^2} \int_{a}^{b}\int_{c}^{d} tsf(t,s)dsdt \right| \leq \frac{(a+b)(c+d)}{16} \left\| \frac{\partial^2 f(t,s)}{\partial t \partial s} \right\|_{\infty}.$$

Lemma 2. *Let $f : [a,b] \times [c,d] \rightarrow \mathbb{R}$ be an absolutely continuous fuction such that the partial derivative of order 2 exist for all $(t,s) \in [a,b] \times [c,d]$. Then, we have*

$$\begin{aligned} f(x,y) &= \frac{1}{m(\alpha_1,\alpha_2)} \int_{\alpha_1}^{\alpha_2} w(t)f(t,y)dt + \frac{1}{m(\beta_1,\beta_2)} \int_{\beta_1}^{\beta_2} w(s)f(x,s)ds \\ &\quad - \frac{1}{m(\alpha_1,\alpha_2)m(\beta_1,\beta_2)} \left[\int_{\alpha_1}^{\alpha_2}\int_{\beta_1}^{\beta_2} w(t)w(s)f(t,s)dsdt - \int_{\alpha_1}^{\alpha_2}\int_{\beta_1}^{\beta_2} P(x,t)Q(y,s)\frac{\partial^2 f(t,s)}{\partial t \partial s}dsdt \right] \end{aligned} \tag{2.13}$$

where

$$P(x,t)=\begin{cases} P_1(\alpha_1,t)=\int\limits_{\alpha_1}^{t} w(u)du, & \alpha_1\le t<x \\ \\ P_2(\alpha_2,t)=\int\limits_{\alpha_2}^{t} w(u)du, & x\le t\le \alpha_2 \end{cases}$$

and

$$Q(y,s)=\begin{cases} Q_1(\beta_1,s)=\int\limits_{\beta_1}^{s} w(u)du, & \beta_1\le s<y \\ \\ Q_2(\beta_2,s)=\int\limits_{\beta_2}^{s} w(u)du, & y\le s\le \beta_2. \end{cases}$$

for all $(\alpha_i,\beta_i)\in[a,b]\times[c,d],\ i=1,2$ *with* $\alpha_1<\alpha_2,\ \beta_1<\beta_2$.

Proof. By definitions of $P(x,t)$ and $Q(y,s)$, we have

$$\int\limits_{\alpha_1}^{\alpha_2}\int\limits_{\beta_1}^{\beta_2} P(x,t)Q(y,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt=\int\limits_{\alpha_1}^{x}\int\limits_{\beta_1}^{y} P_1(\alpha_1,t)Q_1(\beta_1,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt$$

$$+\int\limits_{\alpha_1}^{x}\int\limits_{y}^{\beta_2} P_1(\alpha_1,t)Q_2(\beta_2,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt+\int\limits_{x}^{\alpha_2}\int\limits_{\beta_1}^{y} P_2(\alpha_2,t)Q_1(\beta_1,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt$$

$$+\int\limits_{x}^{\alpha_2}\int\limits_{y}^{\beta_2} P_2(\alpha_2,t)Q_2(\beta_2,s)\frac{\partial^2 f(t,s)}{\partial t\partial s}dsdt.$$

By similar computation in Lemma 1, we get (2.13). □

Theorem 4. *Let* $f:[a,b]\times[c,d]\to\mathbb{R}$ *be an absolutely continuous fuction such that the partial derivative of order* 2 *exist and is bounded, i.e.,*

$$\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty=\sup_{(x,y)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right|<\infty$$

for all $(t,s)\in[a,b]\times[c,d]$. *Then, we have*

$$\left|f(x,y)-\left[\frac{1}{m(\alpha_1,\alpha_2)}\int\limits_{\alpha_1}^{\alpha_2}w(t)f(t,y)dt+\frac{1}{m(\beta_1,\beta_2)}\int\limits_{\beta_1}^{\beta_2}w(s)f(x,s)ds\right]\right.$$

$$\left.-\frac{1}{m(\alpha_1,\alpha_2)m(\beta_1,\beta_2)}\int\limits_{\alpha_1}^{\alpha_2}\int\limits_{\beta_1}^{\beta_2}w(t)w(s)f(t,s)dsdt\right| \tag{2.14}$$

$$\leq \frac{A_1(x)B_1(y)}{m(\alpha_1,\alpha_2)m(\beta_1,\beta_2)}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty$$

where

$$A_1(x)=\int\limits_{\alpha_1}^{x}(x-u)w(u)du+\int\limits_{x}^{\alpha_2}(u-x)w(u)du$$

and

$$B_1(y)=\int\limits_{\beta_1}^{y}(y-u)w(u)du+\int\limits_{y}^{\beta_2}(u-y)w(u)du$$

for all $(\alpha_i,\beta_i)\in[a,b]\times[c,d],\ i=1,2$ *with* $\alpha_1<\alpha_2,\ \beta_1<\beta_2$.

Proof. From Lemma 1, using the properties of modulus and by similar computation in Theorem 3, we get (2.14). The details are omitted. □

Remark 4. *We choose* $\alpha_1=a,\ \alpha_2=b,\ \beta_1=c,\ \beta_2=d$ *in (2.14), then Theorem 4 reduces Theorem 3.*

Corollary 2. *Under the assumptions of Theorem 4, we have*

$$\left|f(\frac{a+b}{2},\frac{c+d}{2})-\left[\frac{1}{m(\alpha_1,\alpha_2)}\int\limits_{\alpha_1}^{\alpha_2}w(t)f(t,\frac{c+d}{2})dt+\frac{1}{m(\beta_1,\beta_2)}\int\limits_{\beta_1}^{\beta_2}w(s)f(\frac{a+b}{2},s)ds\right]\right.$$

$$\left.-\frac{1}{m(\alpha_1,\alpha_2)m(\beta_1,\beta_2)}\int\limits_{\alpha_1}^{\alpha_2}\int\limits_{\beta_1}^{\beta_2}w(t)w(s)f(t,s)dsdt\right|$$

$$\leq \frac{A_1(\frac{a+b}{2})B_1(\frac{c+d}{2})}{m(\alpha_1,\alpha_2)m(\beta_1,\beta_2)}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty$$

where

$$A_1(\frac{a+b}{2})=\int\limits_{\alpha_1}^{\frac{a+b}{2}}(\frac{a+b}{2}-u)w(u)du+\int\limits_{\frac{a+b}{2}}^{\alpha_2}(u-\frac{a+b}{2})w(u)du$$

and

$$B_1(\frac{c+d}{2}) = \int\limits_{\beta_1}^{\frac{c+d}{2}} (\frac{c+d}{2} - u)w(u)du + \int\limits_{\frac{c+d}{2}}^{\beta_2} (u - \frac{c+d}{2})w(u)du.$$

Remark 5. *We choose $w(u) = 1$ in Corollary 2, we get*

$$\left| f(\frac{a+b}{2}, \frac{c+d}{2}) - \left[\frac{1}{(\alpha_2 - \alpha_1)} \int\limits_{\alpha_1}^{\alpha_2} f(t, \frac{c+d}{2})dt + \frac{1}{(\beta_2 - \beta_1)} \int\limits_{\beta_1}^{\beta_2} f(\frac{a+b}{2}, s)ds \right] \right.$$

$$\left. - \frac{1}{(\alpha_2 - \alpha_1)(\beta_2 - \beta_1)} \int\limits_{\alpha_1}^{\alpha_2} \int\limits_{\beta_1}^{\beta_2} f(t, s)dsdt \right|$$

$$\leq \frac{A_3 B_3}{64(\alpha_2 - \alpha_1)(\beta_2 - \beta_1)} \left\| \frac{\partial^2 f(t, s)}{\partial t \partial s} \right\|_\infty$$

where

$$A_3 = (a + b - 2\alpha_1)^2 + (a + b - 2\alpha_2)^2$$

and

$$B_3 = (c + d - 2\beta_1)^2 + (c + d - 2\beta_2)^2.$$

References

[1] N. S. Barnett and S. S. Dragomir, *An Ostrowski type inequality for double integrals and applications for cubature formulae,* Soochow J. Math., 27(1), (2001), 109-114.

[2] S. S. Dragomir, N. S. Barnett and P. Cerone, *An n-dimensional version of Ostrowski's inequality for mappings of Hölder type*, RGMIA Res. Pep. Coll., 2(2), (1999), 169-180.

[3] A. M. Ostrowski, *Über die absolutabweichung einer differentiebaren funktion von ihrem integralmitelwert*, Comment. Math. Helv. 10(1938), 226-227.

[4] B. G. Pachpatte, *On an inequality of Ostrowski type in three independent variables*, J. Math.Anal. Appl., 249(2000), 583-591.

[5] B. G. Pachpatte, *On a new Ostrowski type inequality in two independent variables*, Tamkang J. Math., 32(1), (2001), 45-49

[6] B. G. Pachpatte, *A new Ostrowski type inequality for double integrals*, Soochow J. Math., 32(2), (2006), 317-322.

[7] M. Z. Sarikaya, *On the Ostrowski type integral inequality*, Acta Math. Univ. Comenianae, Vol. LXXIX, 1(2010), pp. 129-134.

[8] N. Ujević, *Some double integral inequalities and applications,* Appl. Math. E-Notes, 7(2007), 93-101.

Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce-TURKEY
E-mail address: `sarikayamz@gmail.com`

Department of Mathematics, Faculty of Science and Arts, Afyon Kocatepe University, Afyon-TURKEY
E-mail address: `hogunmez@aku.edu.tr`